


PEXION® DOSE ADJUSTMENT PROTOCOL

Prepared by the Canine Epilepsy Advisory Group: Sigitas Cizinauskas, Robyn Farquhar, Gualtiero Gandini, Jacques Penderis, Kai Rentmeister, Jean Laurent Thibaud, Sandra Tyler, Holger Volk



Dog on 10 mg/kg twice daily (see dosing table)



*In case of severe seizures, efficacy is seen within 48 hours. See package insert for complete product information.

FLEXIBLE OPTIONS ENABLE EASY DOSING CUSTOMISATION

Pexion® is available in 2 convenient tablet sizes—
scored for more dosing options.

BODY WEIGHT (kg)	NUMBER OF TABLETS (administered twice daily based on the recommended initial dose of 10mg per kg bodyweight)	
	 100 mg tablet	 400 mg tablet
5.0	½	—
5.1–10.0	1	—
10.1–15.0	1½	—
15.1–20.0	—	½
20.1–40.0	—	1
40.1–60.0	—	1½
Over 60	—	2

Assuming that Pexion is well tolerated by the dog, the dose can be increased by 50% to 100% increments up to a maximum dosage of 30mg per kg administered twice daily if necessary.

Australia: Boehringer Ingelheim Pty Limited. ABN 52 000 452 308. Animal Health Division, 78 Waterloo Road, North Ryde NSW 2113. Toll Free 1800 038 037. Fax Number 02 8875 8715.
New Zealand: Boehringer Ingelheim (NZ) Ltd. Animal Health Division, Level 1, Unit 9, 42 Ormiston Road, East Tamaki, Auckland. Toll Free 0800 802 461. Fax Number 09 271 0629.
Restricted Veterinary Medicine. Access is only through a Veterinary Authorisation. Pexion® is a registered trademark of Boehringer Ingelheim Vetmedica GmbH, 55216 Ingelheim/Rhein.
For further information relating to transition, dose adjustment and treatment protocols, please contact our Technical Services Veterinarian on the toll free number.